

ORIGINAL

TSCA NON-CONFIDENTIAL BUSINESS INFORMATION

DOCUMENT DESCRIPTION	DOCUMENT CONTROL NUMBER	DATE RECEIVED
8EHQ-11- 18446	88110000330	9/20/11

COMMENTS:

DOES NOT CONTAIN CBI

RECEIVED
OPPT CBIC

338301



2011 SEP 20 AM 10:49

AkzoNobel

Tomorrow's Answers Today

September 19, 2011

Office of Pollution Prevention and Toxics
EPA East – Room 6428 Attn: Section 8(e)
U.S. Environmental Protection Agency
1201 Constitution Avenue, NW
Washington, DC 20004-3302

**ATTENTION: TSCA 8(e) Notice**

Dear U.S. EPA:

On behalf of Akzo Nobel Surface Chemistry LLC, we are submitting results from an OECD 407, 4-week repeated dose study including a 4-week recovery period on Coco amidopropyldimethylamine (CAS# 68140-01-2) under Section 8(e) of TSCA.

In the OECD 407 study, groups of 5 male and 5 female rats were treated by gavage once daily for up to 29 consecutive days. In this study two satellite groups of 5 male and 5 female rats were included in the control group and the high-dose group. These animals were kept for a 4-week recovery period after treatment for observation of reversibility, persistence or delayed occurrence of systemic / irritative effects. The Control Group was dosed with the vehicle only (corn oil).

The following dose levels were selected as follows:

- Control Group: 0 mg/kg body weight/day
- Low Dose Group: 30 mg/kg body weight/day
- Mid Dose Group: 60 mg/kg body weight/day
- High Dose Group: 120 mg/kg body weight/day

A trend towards a lower body weight gain was noted in treated animals associated with a decrease in food consumption in high-dose group when compared to control group. These effects were reversible during the recovery period.

At necropsy white discoloration of the fore stomach was observed in the mid- and high-dose groups confirming the strong irritative/corrosive properties of the test substance. In the satellite groups no test item-related changes were observed at necropsy. Microscopic examination revealed signs of pronounced irritation of the fore stomach in

**CONTAINS NO CBI**


the mid- and high-dose groups. These signs consisted of acanthosis, hyperkeratosis, erosion/ulcer, edema with mixed infiltrate of inflammatory cell in the submucosa and correlated with the white discoloration seen at necropsy. In the low-dose group changes in the fore stomach were also present but with lower severity and incidence.

Furthermore, the microscopic examination after a 4-week recovery period showed a pronounced decrease in the incidence and severity of changes observed in the fore stomach but the recovery was not complete. Acanthosis and hyperkeratosis were still observed.

The NOAEL with respect to systemic toxic effects was considered to be 60 mg/kg/day because of lower food intake and slight reduction in body weight gain observed in animals treated at 120 mg/kg/day. A NOAEL for local effects could not be established because of irritation observed in the fore stomach of all treated groups. However, in the absence of a similar structure in the human stomach, the relevance of the findings in the rat non-glandular portion of the stomach (fore stomach) with respect to human health is not known.

Please contact me if you have any questions at 312-544-7191.

Sincerely,

A handwritten signature in black ink, reading "Edwin C. Bisinger Jr." in a cursive script.

Edwin C. Bisinger Jr., PhD, DABT
Director, Global Product Safety
HSE Department
Akzo Nobel Services Inc.
525 W. Van Buren St.
Chicago, IL 60607-3835

From: (312) 544-7089
Jennifer Wisniewski

Origin ID: GYYA



525 W Van Buren St
Suite 1800
Chicago, IL 60607

Ship Date: 19SEP11
ActWgt: 0.5 LB
CAD: 103210438/NET3180

Delivery Address Bar Code



SHIP TO: (202) 564-8940

BILL SENDER

Office of Pollution Prevent & Toxic
U.S. Environmental Protection Agency
1201 CONSTITUTION AVE NW
ATTN:SECTION 8(E); EPA EAST-RM 6428
WASHINGTON, DC 20460

Ref # Bisinger 7201-SH526
Invoice #
PO #
Dept #


TUE - 20 SEP A1
STANDARD OVERNIGHT

TRK# 7952 0184 4241

8201

XC RDVA

20460
DC-US
DCA



50F02142060F5F4

After printing this label

- 1 Use the 'Print' button on this page to print your label to your laser or inkjet printer
- 2 Fold the printed page along the horizontal line
- 3 Place label in shipping pouch and affix it to your shipment so that the barcode portion of the label can be read and scanned

Use of this system constitutes your agreement to the service conditions in the current FedEx Service Guide, available on fedex.com FedEx will not be responsible for any claim in excess of \$100 per package, whether the result of loss, damage, delay, non-delivery, misdelivery, or misinformation, unless you declare a higher value, pay an additional charge, document your actual loss and file a timely claim. Limitations found in the current FedEx Service Guide apply. Your right to recover from FedEx for any loss, including intrinsic value of the package, loss of sales, income interest, profit, attorney's fees, costs, and other forms of damage whether direct, incidental, consequential, or special is limited to the greater of \$100 or the authorized declared value. Recovery cannot exceed actual documented loss. Maximum for items of extraordinary value is \$500, e.g. jewelry, precious metals, negotiable instruments and other items listed in our ServiceGuide. Written claims must be filed within strict time limits, see current FedEx Service Guide.